



KAMIYA BIOMEDICAL COMPANY

Caspase-1 Inhibitor Drug Screening Kit

For the effective screening of caspase-1 inhibitors using a fluorometric method

Cat. No. KT-133

For research use only, not for use in diagnostic procedures.

PRODUCT INFORMATION

Caspase-1 Inhibitor Drug Screening Kit **Cat. No. KT-133** **100 tests**

PRODUCT

The **K-ASSAY®** Caspase-1 Inhibitor Drug Screening kit is a convenient fluorometric assay to screen caspase-1 inhibitor samples.

PRINCIPLE

Caspases have been shown to play a crucial role in apoptosis induced by various deleterious and physiologic stimuli. Inhibition of caspases can delay apoptosis, implicating a potential role in drug screening efforts. The Caspase-1 Inhibitor Drug Screening Kit provides an effective means for screening caspase inhibitors using fluorometric methods. The assay utilizes synthetic peptide substrate YVAD-AFC (AFC, 7-amino-4-trifluoromethyl coumarin). Active caspase-1 cleaves the synthetic substrate to release free AFC which can then be quantified by fluorometry. Compounds to be screened can directly be added to the reaction and the level of inhibition of caspase-1 activity can be determined by comparison of the fluorescence intensity in samples with and without the testing inhibitors. The assay is simple, straightforward, and can be performed directly in microtiter plates. Each kit contains 100 units of active caspase-1, sufficient for screening 100 caspase inhibitor samples. Assay conditions have been optimized to obtain the maximal activity.

COMPONENT

• 2X Reaction Buffer	10 ml
• Caspase Substrate YVAD-AFC (1 mM)	0.5 ml
• DTT (1 M)	100 µl
• Active Caspase-1 (Lyophilized)	100 units
• Caspase Inhibitor, Z-VAD-FMK (2 mM)	10 µl

PROTOCOLS

A. General Considerations & Reagent Preparations

- After thawing, store the 2X Reaction Buffer at 4°C. Aliquot enough 2X Reaction Buffer for the number of assays to be performed. Add DTT to the 2X Reaction Buffer immediately before use (10 mM final concentration: add 10 µl of 1.0 M DTT stock per 1 ml of 2X Reaction Buffer).
- Protect YVAD-AFC from light.
- Reconstitute the Active Caspase-1 in 550 µl 2X Reaction Buffer. Aliquot and immediately store at -70°C.

B. Assay Procedure

1. Prepare testing sample in dH₂O to a final volume of 50 µl/well. Add 5 µl of Active Caspase-1. Mix well.

Prepare a background control by omitting the Active Caspase-1 from the reaction mixture. Prepare a positive inhibition control by adding 1 µl of the Caspase-1 Inhibitor (provided with the kit) instead of your testing inhibitor.

2. Prepare a Master Mix for each assay containing the following:
45 µl 2X Reaction Buffer (containing 10 mM DTT)
5 µl 1 mM YVAD-AFC substrate (50 µM final concentration)

3. Mix well and add 50 μ l of the Master Mix to each well to start the reaction.
4. Incubate at 37°C for 0.5-1 hour.
5. Read samples in a fluorescence plate reader equipped with a 400-nm excitation filter and 505-nm emission filter. Comparison of the fluorescence intensity of the testing samples with samples containing no inhibitors to determine the inhibition efficiency of the testing inhibitors.

STORAGE

Store kit at -20°C (Store 2X Reaction Buffer at 4°C after opening).

FOR RESEARCH USE ONLY

KAMIYA BIOMEDICAL COMPANY

12779 Gateway Drive, Seattle WA 98168
Tel: (206) 575-8068 Fax: (206) 575-8094
Email: LifeScience@k-assay.com
www.k-assay.com