

PRODUCT DATA SHEET

Product: Anti-Stromal Cell Derived Factor 2 (SDF2) pAb

Cat. No.: PC-2050 (200 µL)

Species Reactivity:

Human, mouse and rat. Other species not known.

Ig Isotype:

Rabbit IgG

Immunogen:

Recombinant SDF2 (Ser19~Leu211) expressed in *E. coli*

Format:

500 μg/mL of polyclonal antibody in 0.01M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol. Purified by antigen-specific affinity chromatography followed by Protein A affinity chromatography.

Storage:

Store at 4° C for frequent use. Stored at -20° C in a manual defrost freezer for two years without detectable loss of activity. Avoid repeated freezethaw cycles.

Applications and Suggested Dilutions:

- Western blot: Use at 0.5 5 μg/mL.
- Immunohistochemistry: Use at 5 20 μg/mL.
- Immunocytochemistry: Use at 5 20 μg/mL.

The optimal dilution for a specific application should be determined by the researcher.

Limitations:

For *in vitro* research use only. Not for use in diagnostics or in humans.

Warranty:

No warranties, expressed or implied, are made regarding the use of this product. **KAMIYA BIOMEDICAL COMPANY** is not liable for any damage, personal injury, or economic loss caused by this product.