

# **PRODUCT DATA SHEET**

# Product: Anti-MRP2/cMOAT mAb, clone M<sub>2</sub>III-6

## *Cat. No.:* MC-206 (1 mL) MC-164 (0.5 mL)

### Specificity:

M<sub>2</sub>III-6 reacts with an internal epitope of human MRP2 (ABCC2), a 170-180 kDa transmembrane protein known as the canalicular multi-organic anion transporter (cMOAT), which is absent in humans with the Dubin-Johnson syndrome, an autosomal recessive liver disorder characterized by chronic conjugated hyperbilirubinemia. MRP2 is closely related to the multidrug resistance-related protein MRP. MRP2 is overexpressed in a subset of cisplatin-resistant cell lines. Does not cross-react with human *MDR1*, *MRP1*, *MRP3*, or *MRP5* gene products.

#### Species Reactivity:

Reacts with human and rat MRP2. Others not tested.

#### Ig Isotype:

Mouse IgG<sub>2a</sub>

#### Immunogen:

Bacterial fusion protein of MRP2 containing the 202-amino acid COOH-terminal end of the protein.

#### Format:

0.5 mL or 1 mL Protein G purified monoclonal antibody at 100  $\mu$ g/mL in PBS with protein stabilizer and 0.02% sodium azide.

#### Storage:

Store at 4°C for short term. Aliquot and store at -20°C for long term. Avoid repeated freeze/ thaw cycles.

#### Applications and Suggested Dilutions:

- Flow cytometry: Optimal conditions still to be determined.
- Immunocytochemistry
- Immunohistochemistry: Use frozen sections at a 1:20 dilution. Paraffin-embedded tissues can also be used.
- Western blot: Use at a 1:20-1:50 dilution.

The optimal dilution for a specific application should be determined by the researcher.

#### Limitations:

For *in vitro* research use only. Not for use in diagnostics or in humans.

#### Warranty:

No warranties, expressed or implied, are made regarding the use of this product. KAMIYA BIOMEDICAL COMPANY is not liable for any damage, personal injury, or economic loss caused by this product.